

Is the time lag-luminosity relation of GRBs a consequence of the Amati relation?

M. Hafizi¹ and R. Mochkovitch²

¹Tirana University, Faculty of Natural Sciences, Tirana, Albania

²Institut d'Astrophysique de Paris - UMR 7095 CNRS et Université Pierre et Marie Curie,
98 bis, boulevard Arago, 75014 Paris, France

Preprint online version: February 5, 2008

ABSTRACT

Context. The lag-luminosity relation (LLR) provides a way of estimating GRB luminosity by measuring the spectral lags between different energy bands.

Aims. We want to understand the origin of the LLR and test its validity. This appears especially important if the LLR is to be used as a distance indicator.

Methods. We perform a linear analysis of the lag between two spectral bands. The lag is obtained as the time interval between the maxima of a given pulse in the two bands.

Results. We get a simple expression for the lag, which shows in a very simple way how it is related to the spectral evolution of the burst via the variation of the peak energy and spectral indices. When this expression is coupled to the Amati relation, it leads to a LLR that agrees with the observational results only if the burst's spectral evolution is limited to a decrease in peak energy during pulse decay. However, when the variation of the spectral indices is also taken into account, the predicted LLR differs from the observed one.

Conclusions. We briefly discuss some ways to solve this problem, such as a possible correlation between pulse spikiness and burst luminosity.

Key words. gamma rays: bursts; radiation mechanisms: non thermal

1. Introduction

The problem of the distance to gamma-ray bursts remained unsolved until the discovery of the afterglows by Beppo-SAX. Redshifts are now obtained from optical spectra of the afterglow itself or of the host galaxy when the afterglow has faded away. Using the known redshifts, it became possible to calibrate relations by linking absolute burst outputs (luminosity or total radiated energy) and quantities directly available from the observations in gamma-rays. A Cepheid-like relation between variability and luminosity was proposed, for example, by Reichart et al (2001). More recently, Atteia (2003) used the Amati relation (Amati et al 2002) to introduce “pseudo-redshifts” which could be useful to rapidly identify high- z GRBs from their gamma-ray properties alone. In this paper we concentrate on the time lag-luminosity relation (LLR) discovered by Norris et al (2000). The lags were computed by Norris et al using the burst profiles in BATSE band 1 (20 - 50 keV) and 3 (100 - 300 keV). They find that the time

lag Δt_{13} anticorrelates with burst luminosity and propose the following power law relation

$$L = 1.3 \cdot 10^{53} (\Delta t_{13}/0.01 \text{ s})^{-1.15} \text{ erg.s}^{-1}. \quad (1)$$

The origin of the LLR was then investigated by Kocovski and Liang (2003), Ryde (2005), and Ryde et al (2005), who found that the observed lags are a consequence of the burst's spectral evolution. In this contribution we perform a new analysis of the origin of lags and discuss how the LLR may be linked to the Amati relation and possibly also to the variability-luminosity relation.

2. Count rates in different energy bands

We consider a spectral band $[E_i, E_j]$ and assume a spectrum shape consisting of two smoothly-connected power laws of respective slopes α and β at low and high energy (Band et al, 1993). The count rate $N_{ij}(t)$ in band $[E_i, E_j]$ reads

$$N_{ij}(t) = A(t) \int_{x_i}^{x_j} \mathcal{B}_{\alpha\beta}(x) dx \quad (2)$$

where the function $A(t)$ depends on time alone. The limits of the integral are $x_{i,j} = E_{i,j}(1+z)/E_p(t)$, z being the redshift of the source, $E_p(t)$ the peak energy of the instantaneous spectrum (in source rest frame) and $\mathcal{B}_{\alpha\beta}(x)$ the spectrum shape.

Considering now another spectral band $[E_k, E_l]$, we can relate $N_{kl}(t)$ to $N_{ij}(t)$ in the following way

$$\begin{aligned} N_{kl}(t) &= N_{ij}(t) \times \frac{\int_{x_k}^{x_l} \mathcal{B}_{\alpha\beta}(x) dx}{\int_{x_i}^{x_j} \mathcal{B}_{\alpha\beta}(x) dx} \\ &= N_{ij}(t) \times \mathcal{F}_{ijkl} [E_p(t), \alpha(t), \beta(t)] \end{aligned} \quad (3)$$

where $\mathcal{F}_{ijkl}$ can be seen as the “spectral correction” between bands $[E_i, E_j]$ and $[E_k, E_l]$. We simplify the notation by considering only BATSE bands 1 [20, 50 keV] and 3 [100, 300 keV] so that we have

$$N_3(t) = N_1(t) \times \mathcal{F}_{13} [E_p, \alpha, \beta] \quad (4)$$

with

$$\mathcal{F}_{13} = \frac{\int_{100(1+z)/E_p}^{300(1+z)/E_p} \mathcal{B}_{\alpha\beta}(x) dx}{\int_{20(1+z)/E_p}^{50(1+z)/E_p} \mathcal{B}_{\alpha\beta}(x) dx}. \quad (5)$$

We then assume that a given pulse in the burst profile reaches its maximum at a time t_1 (resp. t_3) in band 1 (resp. 3) and we estimate the lag by the difference

$$\Delta t_{13} = t_1 - t_3. \quad (6)$$

Since in most cases the observed lags are small compared to the pulse duration, we evaluate Δt_{13} from a linear analysis of the pulse shape around t_1 . Using $\left. \frac{dN_1}{dt} \right|_{t=t_1} = \dot{N}_1(t_1) = 0$, we can write

$$N_1(t) \simeq N_1(t_1) + \frac{1}{2} \ddot{N}_1(t_1) (t - t_1)^2, \quad (7)$$

while the spectral correction gives to same order

$$\mathcal{F}_{13}(t) = \mathcal{F}_{13}(t_1) + \dot{\mathcal{F}}_{13}(t_1) (t - t_1) + \frac{1}{2} \ddot{\mathcal{F}}_{13}(t_1) (t - t_1)^2, \quad (8)$$

with $\dot{\mathcal{F}}_{13}(t_1)$ and $\ddot{\mathcal{F}}_{13}(t_1)$ being related to the partial derivatives of $\mathcal{F}_{13}$ with respect to E_p , α , and β . For $\dot{\mathcal{F}}_{13}(t_1)$ we have

$$\dot{\mathcal{F}}_{13}(t_1) = \left. \frac{\partial \mathcal{F}_{13}}{\partial E_p} \right|_{t_1} \dot{E}_p(t_1) + \left. \frac{\partial \mathcal{F}_{13}}{\partial \alpha} \right|_{t_1} \dot{\alpha}(t_1) + \left. \frac{\partial \mathcal{F}_{13}}{\partial \beta} \right|_{t_1} \dot{\beta}(t_1), \quad (9)$$

while $\ddot{\mathcal{F}}_{13}(t_1)$ contains nine terms. We now compute the logarithmic derivative of $N_3(t)$ to the first order in $(t - t_1)$

$$\begin{aligned} \frac{\dot{N}_3(t)}{N_3(t)} &= \frac{\dot{N}_1(t)}{N_1(t)} + \frac{\dot{\mathcal{F}}_{13}(t)}{\mathcal{F}_{13}(t)} = \frac{\dot{N}_1(t_1)}{N_1(t_1)} (t - t_1) \\ &+ \frac{\dot{\mathcal{F}}_{13}(t_1)}{\mathcal{F}_{13}(t_1)} + \left[\frac{\ddot{\mathcal{F}}_{13}(t_1)}{\mathcal{F}_{13}(t_1)} - \left(\frac{\dot{\mathcal{F}}_{13}(t_1)}{\mathcal{F}_{13}(t_1)} \right)^2 \right] (t - t_1). \end{aligned} \quad (10)$$

The bracket contains many terms involving partial derivatives of $\mathcal{F}_{13}$ to the first and second order, but it turns out

that they are essentially negligible for the final numerical results. Solving Eq.(10) to get t_3 , such as $\dot{N}_3(t_3) = 0$, finally yields

$$\frac{\Delta t_{13}}{t_p} \simeq \frac{f_{13,E} \dot{E}_p + f_{13,\alpha} \dot{\alpha} + f_{13,\beta} \dot{\beta}}{C_1} \quad (11)$$

with

$$\begin{aligned} f_{13,X} &= \left. \frac{\partial \text{Log} \mathcal{F}_{13}}{\partial \text{Log} X} \right|_{t_1}, \quad \dot{E}_p = \frac{\dot{E}_p}{E_p} t_p \\ \dot{\alpha} &= \frac{\dot{\alpha}}{\alpha} t_p, \quad \dot{\beta} = \frac{\dot{\beta}}{\beta} t_p \quad \text{and} \quad \frac{C_1}{t_p^2} = \frac{\ddot{N}_1(t_1)}{N_1(t_1)} \end{aligned} \quad (12)$$

where t_p is the characteristic duration of the pulse. For two given spectral bands and an assumed spectral shape, Eq.(11) provides a linear estimate of the lag, which directly shows how it is related to burst spectral evolution via the temporal derivatives of E_p , α and β . The “curvature parameter” $|C_1|$ depends on the pulse shape at maximum, large (resp. small) $|C_1|$ values corresponding to spiky (resp. broad) pulses.

3. The lag luminosity relation

Equation (11) gives the lag between BATSE bands 1 and 3 if the values of E_p , α , β , their time derivatives, and the pulse shape are known at maximum. It will become a LLR if these parameters can be related in some way to the luminosity. The Amati relation (Amati et al, 2002) provides such a link but, in its most studied version, it connects the isotropic energy in gamma-rays to the E_p value of the global spectrum. However, it has been suggested that a similar relation may exist between E_p and the luminosity. Yonetoku et al (2004) find, for example, a relation between the maximum luminosity and the global E_p , while Ghirlanda et al (2005) propose a relation between the values of E_p and the luminosity both taken at pulse maximum

$$E_p = 380 \left(\frac{L}{1.6 \cdot 10^{52} \text{ erg.s}^{-1}} \right)^{0.43} \text{ keV}. \quad (13)$$

If a substantial fraction of bursts satisfy Eq.(13), it will, together with Eq.(11), lead to a LLR that can be compared to the observational data.

3.1. A (too) simple example

In this section we limit the burst’s spectral evolution to the variation of E_p only and write $\dot{\alpha} = \dot{\beta} = 0$. Equation (11) then simply becomes

$$\frac{\Delta t_{13}}{t_p} \simeq \frac{f_{13,E} \dot{E}_p}{C_1} \quad \text{with} \quad f_{13,E} = \left. \frac{\partial \text{Log} \mathcal{F}_{13}}{\partial \text{Log} E_p} \right|_{t_1, \alpha, \beta}. \quad (14)$$

The function $f_{13,E}$ tends towards 0 for the highest and lowest values of E_p , while it is at its maximum for $50 \lesssim E_p/(1+z) \lesssim 100$ keV. This is illustrated in Fig.1 and can be easily understood, since for $E \ll E_p$ (resp. $E \gg$




Fig. 1. The function $f_{13,E}$ given by Eq.(14) plotted as a function of the observed peak energy for $\alpha = -1$ and $\beta = -2.25$; full line: linear scale; dashed line: logarithmic scale.

E_p) the Band function $\mathcal{B}(x)$ behaves as x^α (resp. x^β). Therefore for high E_p values ($E_p/(1+z) \gg 300$ keV)

$$\mathcal{F}_{13} \simeq \frac{300^{1+\alpha} - 100^{1+\alpha}}{50^{1+\alpha} - 20^{1+\alpha}} = C^{st} \quad \text{and} \quad f_{13,E} = 0 \quad (15)$$

a similar result with β replacing α being obtained for low $E_p/(1+z) \ll 20$ keV. At intermediate values, $50 \lesssim E_p/(1+z) \lesssim 100$ keV, and making the rough approximation that $\mathcal{B}(x) \propto x^\alpha$ in band 1 while $\mathcal{B}(x) \propto x^\beta$ in band 3, we obtain

$$\mathcal{F}_{13} \simeq \frac{300^{1+\beta} - 100^{1+\beta}}{50^{1+\alpha} - 20^{1+\alpha}} \left(\frac{E_p}{1+z} \right)^{\alpha-\beta} \quad (16)$$

and therefore

$$f_{13,E} = \alpha - \beta. \quad (17)$$

Finally it can be shown that the shape of $\mathcal{B}(x)$ at small x , $\mathcal{B}(x) = x^\alpha [1 - (2 + \alpha)x]$ leads to a power law behavior for $f_{13,E}$ at large E_p as seen in Fig.1.

If the Amati-like relation (Eq.(13)) is satisfied, high luminosity bursts will have a large E_p and therefore a small lag, while lags will be comparatively large for bursts with E_p in the range $50(1+z) - 100(1+z)$ keV, i.e. $L \sim 4 \cdot 10^{50} (1+z)^{2.3}$ erg.s⁻¹ at pulse maximum. Equation (14) also predicts that, for a given luminosity and spectral evolution, spiky bursts (large $|C_1|$) will have smaller lags than bursts with broad pulses (small $|C_1|$), in agreement with

observations (Hakkila & Giblin 2006). The LLR obtained with this simple model is represented in Fig.2 for different values of the ratio $|\dot{e}_p/C_1|$ and a typical burst redshift $z = 1$. A large (resp. small) ratio corresponds to a faster (resp. slower) spectral evolution or to a broader (resp. spikier) pulse. At high luminosity ($L > 10^{52}$ erg.s⁻¹), the LLR has a power law behavior since

$$\frac{d \text{Log} L}{d \text{Log} \Delta t} = \left(\frac{d \text{Log} L}{d \text{Log} E_p} \right) \left(\frac{d \text{Log} E_p}{d \text{Log} \Delta t} \right) \quad (18)$$

the first factor being given by the Amati relation (Eq.13), while the second results from the spectrum shape. From Eq.(14) we have

$$\frac{d \text{Log} E_p}{d \text{Log} \Delta t} = \frac{d \text{Log} E_p}{d \text{Log} f_{13,E}} \quad (19)$$

which is a constant at high E_p (and hence luminosity) values (see Fig.1). However at luminosities smaller than 10^{52} erg.s⁻¹ the model predicts that the power law behavior of the LLR should break down with the lag passing through a maximum and then decreasing. This clearly contradicts GRB 980425, which has both a very low luminosity and a large lag. But GRB 980425 does not satisfy the Amati relation so that its departure from the LLR is not surprising. One should instead consider that this burst has a broad temporal profile, i.e. a small $|C_1|$ and an E_p of 138 keV (Ghisellini et al 2006) corresponding to the maximum of $f_{13,E}$ (see Fig.1) and therefore to a large expected lag.




Fig. 2. Lag-luminosity relation resulting from Eq.(14) which limits the spectral evolution to a decrease of E_p during pulse decay (α and β being kept constant respectively equal to -1 and -2.25). The three lines cover an order of magnitude in $|\dot{e}_p/C_1|$ from 0.1 (dashed line) to 0.3 (full line) and 1 (dotted line). A redshift value $z = 1$ has been assumed.

3.2. A more complete study

The spectral evolution of GRBs is, however, not limited to a decrease in E_p during pulse decay. A hard-to-soft evolution is also observed for the spectral indices α and β . In some extreme cases α has been seen to decrease from $\alpha \sim 1.5$ (a value a priori excluded by the synchrotron model) to about -0.5 in just a few seconds (Crider et al 1997). When the variation in the spectral indices is included in our linear analysis, it no longer predicts a vanishing lag for high or low E_p and L values since now $f_{13,\alpha} \neq 0$ when $E_p \rightarrow \infty$ and $f_{13,\beta} \neq 0$ when $E_p \rightarrow 0$ (for example $\lim_{E_p \rightarrow \infty} f_{13,\alpha} \simeq -1.7$ for $\alpha = -1$ and $\beta = -2.25$). The lag then reaches a constant limiting value at low and high luminosities, where it apparently contradicts the observed LLR. This is shown in Fig.3 where our calculated LLR has been plotted for different values of $\dot{a}$ and a fixed $\dot{b} = 0.1$. Even a moderate variation in the spectral indices has a dramatic effect on the LLR and the global agreement with the Norris et al (2000) results that was found in the last section is now lost.

4. Discussion

These results clearly disagree with the observational data for the most luminous GRBs. Therefore if real bursts do satisfy the LLR proposed by Norris et al. (2000), a solution

has to be found for the apparent discrepancy between our analysis and the observations:

(i) A first option could be that in most cases the variation in the spectral index α is small, at least around pulse maximum. However this does not seem to be the case for the bright events for which a detailed, time resolved, spectral analysis has been possible (Preece et al 2000). Moreover, the constraint on any variation in α appears so severe (only the LLR with $\dot{a} = 0.01$ in Fig.3 is marginally compatible with the Norris et al. results) that it seems difficult to expect it will be satisfied by a large fraction of GRBs.

(ii) A more interesting possibility would be that a relation may exist between the curvature parameter $|C_1|$ and the luminosity, bursts with spiky pulses being more luminous than bursts with broad pulses. This might be a different way to express the variability-luminosity relation proposed by Reichart et al (2001). We tried, for example, a simple linear expression of the form

$$|C_1| = 2 + 0.2(L_{51} - 1) \quad (20)$$

where the resulting LLR with $|\dot{e}_p/C_1| = 0.3$ and $\dot{a} = \dot{b} = 0.1$ is represented in Fig.3. In spite of the variation in α , it now gives again very small lags at high luminosity because the pulses are then much spikier than at low luminosity.




Fig. 3. LLR with $|\dot{e}_p/C_1| = 0.3$, but now also including the variation in the low and high-energy spectral indices. The thin lines correspond respectively (from left to right) to $\dot{a} = 0.01, 0.03, 0.1$, and 0.3 (and have all $\dot{b} = 0.1$), while the thick full line represents the $\dot{a} = \dot{b} = 0$ case. The dashed line is obtained with $\dot{a} = \dot{b} = 0.1$ and a varying curvature parameter given by Eq.(20).

5. Conclusion

We have performed a linear analysis of the time lag between two spectral bands and have obtained a simple relation (Eq.(11)) which clarifies how the lag is related to the burst spectral evolution. When this relation is used in conjunction with the Amati relation it leads to a satisfactory LLR only if the spectral evolution of GRBs is limited to a decrease of E_p during pulse decay. If the variation in the spectral indices is also included, the lag does not decrease any longer to low values, even at very high burst luminosity. We have briefly discussed the possibility that short lags might be recovered if burst luminosity is correlated to the shape of the pulses, bursts with spiky pulses being more luminous than bursts with broad pulses. The available sample of GRBs with both measured lags and known distance is still small but should increase with SWIFT. This will allow constraining tests of the results presented in this paper.

References

- Amati, L., Frontera, F., Tavani, M., et al. 2002, A&A 390, 81.
- Atteia, J.L. 2003, A&A 407, L1.
- Band, D., Mateson, J., Ford, L., et al. 1993, ApJ, 413, 281.
- Crider, A. et al 1997, ApJ 479, L39.
- Ghirlanda, G., Ghisellini, G. & Firmani, C., et al. 2005, MNRAS 360, L45.
- Ghisellini, G., Ghirlanda, G., Mereghetti, S. et al. 2006, astro-ph/0605431.
- Hakkila, J. & Giblin, T.W. 2006, astro-ph/0604151
- Kocevski, D. & Liang, E. 2003, ApJ 594, 385.
- Norris, J., Marani G. & Bonnell, J. 2000, ApJ 534, 248.
- Preece, R.D., Briggs, M.S., Mallozi, R.S., et al. 2000, ApJS 126, 19.
- Reichart, D., Lamb, D., Fenimore, E., et al 2001, ApJ 552, 57.
- Ryde, F. 2005, A&A 429, 869.
- Ryde, F., Kocevski, D, Bagoly, Z, et al 2005, A&A 432, 105.
- Yonetoku, R. et al 2004, ApJ 609, 935.

Acknowledgements. The authors thank Frédéric Daigne for helpful discussions and Felix Ryde for communicating several of his results prior to publication. They also thank the anonymous referee whose comments have considerably improved an initial version of this paper.